

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

A4

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>5</sup> :</b> <b>A61K 31/28, C07F 17/00, 7/28</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 94/04142</b> <b>(43) International Publication Date:</b> 3 March 1994 (03.03.94)
<b>(21) International Application Number:</b> PCT/US93/07875 <b>(22) International Filing Date:</b> 19 August 1993 (19.08.93)  <b>(30) Priority data:</b> 102866 19 August 1992 (19.08.92) IL  <b>(71)(72) Applicant and Inventor:</b> KEINAN, Ehud [IL/US]; 6549 Dennison Avenue, San Diego, CA 92122 (US).  <b>(74) Agent:</b> DIPPert, William, H.; Cowan, Liebowitz & Lat- man, 605 Third Avenue, New York, NY 10158 (US).  <b>(81) Designated States:</b> AT, AU, BB, BG, BR, CA, CH, CZ, DE, DK, ES, FI, GB, HU, JP, KP, KR, LK, LU, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SK, UA, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI pa- tent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).		<b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> NOVEL METALLOCENES AS ANTI-TUMOR DRUGS  <b>(57) Abstract</b>  The invention relates to novel titanocene derivatives possessing chemotherapeutic activity and method for their prepara- tion. These compounds possess two cyclopentadiene rings linked to titanium as a central atom and bound covalently to two phen- oxy groups which possess a substituent R selected from the group consisting of: COOCH <sub>3</sub> , COOC <sub>2</sub> H <sub>5</sub> , H, COOCH <sub>2</sub> CH <sub>2</sub> OCH <sub>2</sub> CH <sub>2</sub> OCH <sub>3</sub> and are free from amino groups, nitro, chloride and fluoride. The novel compounds repres- ent a compromise between the main properties for an antitumor agent, i.e. electrophilicity and stability, being water soluble. Cytotoxicity measurements of these compounds showed significant growth inhibition properties, expressed in terms of IC <sub>50</sub> [M] values.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NE	Niger
BE	Belgium	CN	Guinea	NL	Netherlands
BF	Burkina Faso	GR	Greece	NO	Norway
BG	Bulgaria	HU	Hungary	NZ	New Zealand
BJ	Benin	IE	Ireland	PL	Poland
BR	Brazil	IT	Italy	PT	Portugal
BY	Belarus	JP	Japan	RO	Romania
CA	Canada	KP	Democratic People's Republic of Korea	RU	Russian Federation
CF	Central African Republic	KR	Republic of Korea	SD	Sudan
CG	Congo	KZ	Kazakhstan	SE	Sweden
CH	Switzerland	LJ	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovak Republic
CM	Cameroon	LU	Luxembourg	SN	Senegal
CN	China	LV	Latvia	TD	Chad
CS	Czechoslovakia	MC	Monaco	TG	Togo
CZ	Czech Republic	MG	Madagascar	UA	Ukraine
DE	Germany	ML	Mali	US	United States of America
DK	Denmark	MN	Mongolia	UZ	Uzbekistan
ES	Spain			VN	Viet Nam
FI	Finland				

## NOVEL METALLOCENES AS ANTI-TUMOR DRUGS

The present invention relates to new titanocene compounds. More particularly the invention relates to new titanocene complexes and methods for their preparation, which possess chemotherapeutic activity being useful for the treatment of human tumors.

BACKGROUND OF THE INVENTION

There are known metallocene complexes containing titanium, vanadium, niobium and molybdenum as a metal ion, which are active against a variety of tumor cell lines such as B16 melanoma, colon 38 carcinoma, Lewis lung carcinoma, etc. It has been shown that the activity of vanadium complexes related to the formula  $Cp_2VCl_2$  where  $Cp$  is cyclopentadiene, against human epidermoid (HEP-2) tumor cells in vitro and against mouse tumor cells, is similar to that of cis-platin (Murthy M.S. et al. Proc. Am. Assoc. Cancer Res. 1986, 27, 279). A study which was carried out with a corresponding molybdenum compound, supports the possibility that these complexes are binding 5'-phosphate terminated polynucleotides, thus inhibiting DNA replication, by a mechanism which is different from that of cis-platinum complexes (Kon, L.Y. et al. J. Am. Chem. Soc. 1991, 113, 9027).

Titanocene dichloride, one of the first metallocene compounds which was tested, was found to be indeed a very reactive anti-tumor reagent. Due to its rapid hydrolysis

to the corresponding dihydroxy derivative, it is quite reasonable to assume that this dihydroxy titanocene is the actual drug. Accordingly, many references can be found describing titanocene compounds which were tested  
5 in an attempt to possess an improved cytotoxicity. Examples of such compounds include halides, pseudohalides, carboxylates, and phenolates. However, no significant improvement over titanocene dichloride in the antitumor activity has been achieved.

10 The metallocene diacido complexes, having the general formula  $(C_5H_5)_2MX_2$  are characterized by the following structural features:

- The geometry of the complexes is that of a distorted tetrahedron.
- 15 - The complexes contain two uninegative acido ligands X coordinated to the central metal atom and arranged in adjacent "cis-like" position.
- The sites of the other two ligands are occupied by two anionic cyclopentadienyl rings.
- 20 Attempts to modify the cyclopentadienide rings lead to a decreased biological activity.

In a very recent U.S. Patent No. 5,002,969 there are described cytostatic pharmaceutical compositions based on titanocene complexes. A group which is present in all  
25 these complexes is an amino or substituted amino bound to

the titancceno moiety. These compounds are obtained by a reaction between a titanocene dihalogenide and an amino phenol, lithium aminophenolate, or lithium amino thiophenolate. There is mentioned that the compounds have a better solubility in water than titanocene dichloride, fact which improves their application and dosing.

Other titanocene complexes which were described, differ by their ionic character from the neutral titanocene compounds. Most of them correspond to the general formula  $[(C_5H_5)_2TiXL]^+Y^-$  where X and Y are anions and L is a neutral donor molecule. These ionic titanocene complexes are characterized by their improved water solubility compared with the neutral titanocene compounds.

It is an object of the present invention to provide novel titanocene derivatives. It is another object of the present invention to provide novel titanocene derivatives which possess a superior cytotoxic activity than the cis-platinum complexes.

#### BRIEF DESCRIPTION OF THE INVENTION.

The invention relates to novel titanocene derivatives which comprise two cyclopentadiene rings linked to titanium as a central atom, which are bound covalently to two phenoxy groups which possess a substituent R which is selected from the group consisting of:  $COOCH_3$ ,  $COOC_2H_5$ , H,  $COOCH_2CH_2OCH_2CH_2OCH_3$  being free from amino groups,

nitro group, chloride and fluoride. The above novel titanocene derivatives represent a compromise between the two main properties required for an antitumor agent: electrophilicity and stability.

5

DETAILED DESCRIPTION OF THE INVENTION.

It is a generally accepted assumption that titanocenes, as well as other antitumor agents, do react with DNA in a similar manner. Therefore, the two main properties required for the drugs, in addition to the water solubility, are electrophilicity and stability in order to survive the aqueous biological medium during the time required to reach the target. The inventor's approach was to synthesize the new compounds which should possess these two main properties. Accordingly, the titanocene compounds envisaged should contain groups such as phenolates, having the role of moderate leaving groups, and appropriate substituents on the phenyl rings which impart stability to these compounds. Thus, considering the electrophilic role played by the metallocene drug in binding to the nucleophilic sites of polynucleotides, it may be concluded that an optimal biological activity would be achieved when the titanocene compounds, according to the present invention, will contain leaving groups of moderate reactivity, such as phenols substituted at their 4-position with  $\text{COOCH}_3$ ,  $\text{CO}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{OCH}_2\text{CH}_2\text{OCH}_3$ ,  $\text{CH}_2\text{-CH}_3$ ,

25

COCH<sub>3</sub>, H and of course possessing a satisfactory hydrolytic stability.

Typical examples of the novel titanocene derivatives are as follows:

- 5 1. Bis(4-cyanophenolato)bis(n<sup>5</sup>-cyclopentadienyl)titanium (IV), hereinafter referred to as TCN.
2. Bis(4-methoxycarbonylphenolato)bis(n<sup>5</sup>-cyclopentadienyl)titanium(IV), hereinafter referred to as TPE
3. Bis(4-ethoxycarbonylphenolato)bis(n<sup>5</sup>-cyclopentadienyl) hereinafter referred to as TEE1.
- 10 4. Bis[4-(2-methoxyethoxy)ethoxy]carbonylphenolato]bis-(n<sup>5</sup>-cyclopentadienyl)titanium(IV), hereinafter referred to as TEG.
5. Bis[4-(methoxy)ethoxycarbonylphenolato]bis(n<sup>5</sup>cyclopentadienyl)titanium(IV), hereinafter referred to as
- 15 TMEM.
6. Bis[4-(2-dimethylamino)ethoxycarbonylphenolato]bis-(n<sup>5</sup>-cyclopentadienyl)titanium(IV), hereinafter referred to as TCA.
- 20 7. Bis[4-(2-trimethylammonio)ethoxycarbonylphenolato]bis-(n<sup>5</sup>cyclopentadienyl)titanium(IV), hereinafter referred to as TCE.

Cytotoxicity measurements carried out with the above compounds show significant growth inhibition properties

25 of these compounds expressed in terms of IC<sub>50</sub>[M] values.

In the following Table 1 are presented the results which show that these compounds are much superior than the known titanocene dichloride (TDC) under the same conditions. The value of the ratio Ti/Pt represents the relative activity of TPE as compared with that of cis-platinum. The first four entries represent data of normal cell lines and the other ten entries represent the experiments with tumor cell lines.



TABLE 1. Cytotoxic data of titanocene derivatives.

	Cell line	Cell type	TPE	TDC	cisPt	Ti Pt
5	CHO	Chinese Hamstead Ovary	$1.3 \times 10^{-5}$	$10^{-3}$	$3.1 \times 10^{-5}$	2
	HMEC	Normal Human Mammary	$3.1 \times 10^{-6}$	$1.3 \times 10^{-4}$	$6.3 \times 10^{-5}$	20
	NHDF	Normal Human Skin	$1.6 \times 10^{-6}$	$10^{-3}$	$3.1 \times 10^{-5}$	20
10	NHEK	Normal Keratino Epithelial	$3.1 \times 10^{-6}$	$10^{-3}$	$6.3 \times 10^{-4}$	200
	Capan 1	Pancreas Carcinoma	$3.9 \times 10^{-7}$	$5.0 \times 10^{-4}$	$3.9 \times 10^{-6}$	10
15	HT-29	Colon Carcinoma	$3.9 \times 10^{-4}$	$5.0 \times 10^{-4}$	$1.3 \times 10^{-4}$	200
	SK-Mel-28	Melanoma	$1.6 \times 10^{-6}$	$10^{-3}$	$1.6 \times 10^{-5}$	10
	H-322	Lung Carcinoma	$6.3 \times 10^{-7}$	$10^{-3}$	$6.3 \times 10^{-5}$	100
	UCLA-P3	Lung Carcinoma	$3.1 \times 10^{-6}$	$10^{-3}$	$6.3 \times 10^{-5}$	20
	MCF-7	Breast Cancer	$2.0 \times 10^{-6}$	$10^{-3}$	$1.3 \times 10^{-4}$	100
20	HL-60	B-cell Leukemia	$3.1 \times 10^{-6}$	$10^{-3}$	$7.8 \times 10^{-6}$	2
	Molt-4	T-cell Leukemia	$10^{-6}$	$10^{-3}$	$10^{-5}$	10
	Ovcar-3	Ovarian Carcinoma	$6.3 \times 10^{-6}$	$10^{-3}$	$6.3 \times 10^{-5}$	10
	P-388	Mouse Leukemia	$3.9 \times 10^{-6}$	$5.0 \times 10^{-4}$	$9.8 \times 10^{-7}$	025

The cytotoxicity results with a number of titanocene derivatives, expressed in concentrations (M) are presented in the attached Table 2 for a number of solid tumors. For combating solid tumors, the titanocene derivatives according to the present invention may be employed as such or as pharmaceutical compositions containing at least one titanocene complex as described above in addition to pharmaceutically acceptable excipients, diluents and/or auxiliary agents. The excipient can serve as an agent for promoting absorption of the medicament by the body or as formulation auxiliary, sweetener, flavouring agent, colourant or preservative. The pharmaceutical formulations of the active compounds are preferably in the form of unit doses matched to the particular mode of administration. The amount of the active compound is chosen so that one or more units are usually sufficient for an individual therapeutic administration. In addition to that, the medicaments with the active compound, may contain also one or more other pharmacologically active constituents, such as: alkylating agents, antimetabolites antibiotics, vitamins, enzymes and heavy metal compounds. The novel titanocene derivatives, according to the present invention, can be prepared from common chemical reagents using standard equipment. It should be realized, that the Examples for their preparations presented

hereinafter are only for illustration and many other routes may be conceived for their syntheses.

**EXAMPLE 1. Preparation of Bis(4-cyanophenolato)bis**  
**(n<sup>5</sup>-cyclopentadienyl)titanium(IV) TCN.**

5 An amount of 238 mg(2mmol) of 4-cyanophenol was dissolved in 10 ml of benzene and 200 mg of sodium hydride 80% in mineral oil (6.67mmol) was added and stirred at room temperature for about 10 minutes. To this mixture an amount of 249 mg (1mmol) of titanocene dichloride was  
10 added and the mixture refluxed for 8 hours, cooled to room temperature and placed on a short column containing silica gel (pre-washed with acetone). The elution with methylene chloride followed by removal of the solvent under reduced pressure, yielded crude TCN. By purifying  
15 the crude TCN on a chromatographic column (silica gel, ethyl acetate-hexane), an amount of 290 mg of pure TCN (70% yield) was obtained in the form of a yellow solid. The analysis of the product on <sup>1</sup>H NMR (CDCl<sub>3</sub>) was as follows:

20 7.52 (d, J=8.6Hz, 4H), 6.64(d, J=8.6Hz, 4H), 6.31(s, 10H).

**EXAMPLE 2. Preparation of Bis(methoxycarbonylphenolato)**  
**bis(n<sup>5</sup>-cyclopentadienyl)titanium(IV) TPE.**

In the same manner as in Example 1, an amount of 273mg (2 mmol) of methyl 4-hydroxybenzoate was reacted with 249 mg

(1mmol) of titanocene dichloride. An amount of 364 mg of TPE (81% yield) in the form of a yellow solid was obtained.

The analysis of the product on  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ) was as follows:

7.91 (d,  $J=8.7$  Hz, 4H), 6.64 (d,  $J=8.7$  Hz, 4H), 6.33 (s, 10H), 2.56 (s, 6H).

**EXAMPLE 3. Preparation of Bis(4-ethoxycarbonylphenolato)-bis( $n^5$ -cyclopentadienyl)titanium(IV) TEE1.**

In the same manner as in Example 1, an amount of 332mg (2mmol) of ethyl 4-hydroxybenzoate was reacted with 249mg (1 mmol) of titanocene dichloride. An amount of 417 mg of TEE1 (81% yield) was obtained.

The analysis of the product on  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ) was as follows:

7.90 (d,  $J=8.6$  Hz, 4H), 6.61 (d,  $J=8.6$  Hz, 4H), 6.31 (s, 10H), 2.94 (q,  $J=7.3$  Hz, 4H), 1.21 (t,  $J=7.3$  Hz, 6H).

**EXAMPLE 4. Preparation of Bis[4-(2-(2-methoxyethoxy)ethoxy]carbonylphenolato]bis( $n^5$ -cyclopentadienyl)titanium(IV) TEG**

(a) In a first step, an amount of 1g (43 mmol) of sodium was dissolved in 25 ml of 2-(2-methoxyethoxy)ethanol. To the resulted solution an amount of 3 g (22mmol) of methyl 4-hydroxybenzoate was added and the mixture was heated to

130°C for 24 hours; after cooling to room temperature, it was acidified with a hydrochloric acid solution (3N) and extracted with ethyl acetate. The removal of the solvent under reduced pressure and column chromatography of the residue (silica gel, hexane:ethyl acetate 3:1) afforded  
5 2-(2-methoxyethoxy)ethyl 4-hydroxybenzoate, in the form of a colourless oil in essentially quantitative yield.

(b) In the second step, an amount of 480 mg (2mmol) of the product obtained in step (a), was reacted with 249 mg  
10 (1mmol) of titanocene dichloride, as described in Example 1. An amount of 355 mg of TEG (54% yield) was obtained. The analysis of the product on <sup>1</sup>H NMR (CDCl<sub>3</sub>) was as follows:

7.95 (d, J=8.6 Hz, 4H), 6.60 (d, J=8.6 Hz, 4H)  
15 6.31 (s, 10H), 4.45 (t, J=5.0Hz, 4H),  
3.83 (t, J=5.0Hz, 4H), 3.70 (t, J=4.6 Hz, 4H),  
3.57 (t, J=4.6 Hz, 4H), 3.38(s, 6H).

**EXAMPLE 5: Preparation of Bis[4-(2-methoxy)ethoxycarbonyl-phenolato]bis(n<sup>5</sup>-cyclopentadienyl)titanium(IV) TMEM.**

20 (a) In the first step (as in the Example 4) 1 g (43 mmol) of sodium was dissolved in 25 ml of 2-methoxyethanol. An amount of 3.0 g (22 mmol) of methyl 4-hydroxy-benzoate was added, producing 2-methoxyethyl 4-hydroxybenzoate, as a colourless oil, in essentially quantitative yield.

(b) In the second step, an amount of 392 mg (2mmol) of the product obtained in step (a) was reacted with 249 mg (1.1 mmol) of titanocene dichloride as described above in Example 1, affording 330 mg of TMEM (58% yield).

5 The analysis of the product on  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ) was as follows:

7.96 (d,  $J=8.6$  Hz, 4H), 6.60 (d,  $J=8.6$  Hz, 4H),  
4H), 6.29 (s, 10H), 4.42 (t,  $J=4.8$  Hz, 4H),  
3.70 (t,  $J=4.8$  Hz, 4H), 3.70 (t,  $J=4.8$  Hz, 4H),  
10 3.40 (s, 6H).

**EXAMPLE 6: Preparation of Bis[4-(2-dimethylamino)ethoxy-  
carbonylphenolato]bis( $n^5$ -cyclopentadienyl)  
titanium(IV) TCA**

(a) In a first step, 1 g (43mmol) of sodium was dissolved  
15 in 20 ml of 2-(dimethylamino)ethanol. An amount of 3.0 g  
(22 mmol) of methyl 4-hydroxybenzoate was added and heated to  $110^\circ\text{C}$  for 24 hours and then cooled to room temperature. The solvent was removed under reduced pressure and using a column chromatography (silica gel, chloroform -  
20 methanol), a white solid of 2-(dimethylamino)-ethyl 4-hydroxybenzoate was obtained.

(b) In the second step, an amount of 418 mg (2 mmol) of the product obtained in step (a) was reacted with 249 mg (1mmol) of titanocene dichloride, as described in Example  
25 1, affording 330 mg of TCA (58% yield).

The analysis of the product on  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ) was as follows:

7.94 (d,  $J=8.6$  Hz, 4H), 6.59 (d,  $J=8.6$  Hz, 4H),

6.30 (s, 10H), 4.39 (t,  $J=7.2$  Hz, 4H),

5 2.71 (t,  $J=7.2$  Hz, 4H), 2.34 (s, 12H).

**EXAMPLE 7: Preparation of Bis[4-(2-trimethylamino)ethoxy-carbonylphenolato]bis( $n^5$ -cyclopentadienyl) titanium(IV).**

10 The TCA product as obtained in the previous Example 6, was treated with an excess of methyl iodide (10 equiv) in benzene for about 6 hours. A yellow solid of TCE is formed, collected by filtration, washed by benzene and ether and dried.

15 The analysis of the product on  $^1\text{H}$  NMR (DMSO) was as follows:

7.87 (d,  $J=8.6$  Hz, 4H), 6.66 (d,  $J=8.6$  Hz, 4H),

6.44 (s, 10H), 4.64 (m, 4H), 3.77 (m, 4H), 3.18 (s, 18H).

-14-

TABLE 2: DATA ON CYTOTOXICITY (log).

Column 1	Column 2	cis- platin	TEG	TMEM	TEE-1	TCE	TPE	TP
1	CHOChin.							
	Harm.Ov.	-4.51		-4.30	-4.60		-4.80	-4.20
2	HMEC Nor.Hum.	-4.20		-4.60	-6.11		-5.70	-3.89
3	NHDF Nor.Humn Skin	-4.51		-4.30	-4.89		-5.41	-4.51
4	Capan-1 Pan (a)	-5.41		-4.89	-5.51	-4.00	-6.41	-4.20
5	HT-29 Colon car.	-3.89		-4.60	-5.20	-4.00	-5.41	-4.20
6	SK ML-28 Melanoam	-4.80		-4.60	-5.51		-5.41	-3.89
7	H-322 Lung Car.	-4.20		-4.89	-5.80	-4.00	-6.20	-4.20
8	UCLA-P3 Lung Car.	-4.20			-5.20		-5.51	-3.89
9	MCF-7 Mamary Car.	-3.89		-4.89	-5.80	-4.30	-5.70	-4.20
10	HL-60 Leukemia	-5.11		-4.60	-5.51		-5.70	-3.30
11	Molt-4 Leukemia	-5.00	-4.00	-7.00	-7.00	-4.00	-6.00	-4.00
12	P-388 Mouse	-6.01			-4.89		-5.41	-4.20
13	NHEK Normal	-3.20			-5.80		-5.51	
14	Ovar-3 Ovarian car.	-4.20	-4.00	-4.60	-5.20	-4.00	-5.20	-3.89
15	SIHA Cerv. Carc.			-4.60		-4.00		
16	MCF-7 (Adr) Adri (b)			-5.51				
1	-3.00 -3.00	-3.00	-3.00		-4.00	-4.60	-3.00	-4.30
2	-3.30 -3.00	-3.00	-3.00		-4.60	-5.20	-3.89	-4.00
3	-3.30 -3.00	-3.00	-3.00		-4.00	-4.00	-3.00	-4.00
4	-3.60 -3.00	-3.00	-3.00		-4.30	-5.51	-3.30	-4.30
5	-3.60 -3.30	-3.30	-3.00		-4.00	-4.89	-3.30	-4.30
6	-3.30 -3.00	-3.00	-3.00		-4.60	-5.51	-3.00	-4.00
7	-3.30 -3.00	-3.00	-3.00	-4.00	-4.30	-4.30	-3.00	-4.30
8	-3.60 -3.00	-3.00	-3.00	-4.30			-3.00	
9	-3.30 -3.00	-3.00	-3.00	-4.30	-4.89	-4.89	-3.00	-4.30
10	-3.00 -3.60	-3.00	-3.00	-4.00	-4.00	-5.80	-3.00	-4.00
11	-2.00 -3.00	-3.00	-3.00		-4.00	-4.00	-3.00	-4.00
12	-3.60 -3.00	-3.00	-3.00	-5.00			-3.30	
13							-3.00	
14	-3.60 -3.00	-3.00	-3.00		-4.30	-5.51	-3.00	-4.00
15					-4.30	-4.30		-4.30
16								

(a) Pancrease Car.

(b) Adriyamicin.

SUBSTITUTE SHEET



C L A I M S :-

1. Novel titanocene derivatives which comprise two cyclopentadiene rings linked to titanium as a central atom, which are bound covalently to two phenoxy groups which possess a radical substituent R which is selected from the group consisting of:

H, COOCH<sub>3</sub>, COOC<sub>2</sub>H<sub>5</sub>, COOCH<sub>2</sub>CH<sub>2</sub>OCH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>,

being free from amino group, nitro group, chloride and fluoride.

2. The novel titanocene derivatives according to Claim 1, which possess chemotherapeutic activity being used for the treatment of tumors.

3. Bis(4-cyanophenolato)bis(n<sup>5</sup>-cyclopentadienyl)titanium (IV).

4. Bis(4-methoxycarbonylphenolato)bis(n<sup>5</sup>-cyclopentadienyl)titanium(IV).

5. Bis(4-ethoxycarbonylphenolato)bis(n<sup>5</sup>-cyclopentadienyl)titanium(IV).

6. Bis(4-[2-(methoxyethoxy)ethoxy]carbonylphenolato)bis(n<sup>5</sup>-cyclopentadienyl)titanium(IV).

7. Bis[4-(2-methoxy)ethoxycarbonylphenolato]bis( $\eta^5$ -cyclopentadienyl)titanium(IV).
8. Bis[4-(2-dimethylamino)ethoxycarbonylphenolato]bis( $\eta^5$ -cyclopentadienyl)titanium(IV).
9. Bis[4-(2-trimethylamino)ethoxycarbonylphenolato]bis( $\eta^5$ -cyclopentadienyl)titanium(IV).
10. The novel titanocene derivatives according to Claim 1, to be applied as medicaments in combination with other pharmaceutically active constituents.

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US93/07875

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(5) : A61K 31/28; C07F 17/00, 7/28

US CL : 556/53, 55; 514/492

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 556/53, 55; 514/492

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

N/A

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

REGISTRY DATA BASE

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Journal of Organometallic Chemistry, Volume 11, No. 3, issued March 1968, K. Andra, "Dicyclopentadienylyltitan-diphenoxide" pages 567-570. see page 567, especially table I, compound (II).	1
A	US, A, 5,002,969 (Kopf-Maier et al) 26 March 1991, see entire document.	1-10
Y	R. Feld et al., "The Organic Chemistry of Titanium" published 1965 by Butterworths Inc. (Washington, D.C.), see pages 3-15, especially page 5, reaction equation No. 1.13.	1

<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.	
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be part of particular relevance "E" earlier document published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "Z" document member of the same patent family
Date of the actual completion of the international search  18 NOVEMBER 1993	Date of mailing of the international search report  06 DEC 1993
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. NOT APPLICABLE	Authorized officer  PORFIRIO NAZARIO-GONZALEZ Telephone No. (703) 308-1235

Form PCT/ISA/210 (second sheet)(July 1992)\*

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP90/07875

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	P.C. Wailes et al., "Organometallic Chemistry of Titanium, Zirconium, and Hafnium", published 1974 by Academic Press (N.Y.), see pages 62-73.	1-10

Form PCT/ISA/210 (continuation of second sheet)(July 1992)\*